

**THE ROLE OF SUPER-PARAMETERIZATION
IN THE CLOUD PARAMETERIZATION PROBLEM**

AKIO ARAKAWA

**A PRELIMINARY TEST OF SUPER-PARAMETERIZATION
IN AN IDEALIZED FRAMEWORK□**

JOON-HEE JUNG and AKIO ARAKAWA

Historical Assessment

Arakawa, A., 2003: The Cumulus Parameterization Problem: Past, Present, Future
(To be submitted to JC)

- GCMs that use cumulus parameterizations constructed with quite different rationale can still produce "comparable" results.
- Thus, cumulus parameterization did not seem to be too demanding problem.
- All surviving cumulus parameterizations can do the simple basic task:
Providing a negative feedback to large-scale destabilization.
- This negative feedback tends to hide model differences and deficiencies, especially when SST is fixed.

**We cannot continue to rely on this kind of "luck"
in future climate models
as the objectives of cumulus parameterization expand.**

OBJECTIVES OF CUMULUS PARAMETERIZATION

Classical Objective

Vertical distributions of cumulus heating (cooling) and cumulus drying (moistening).

Nonclassical Objectives

- (i) Generation, transport and mixing of liquid and ice phases
- (ii) Interactions with the subcloud layer
- (iii) Interactions with radiation
- (iv) Mechanical interactions with the mean flow




Unified cloud (or even unified physics) parameterization

DILEMMA

- Unified cloud parameterization is an extremely challenging scientific problem of scale interactions in the moist atmosphere.
- It is a crucially important engineering problem in weather and climate predictions, for which quick answers are needed.

The struggle in our mind
between the scientific and engineering aspects
is at least partially responsible for
the slow progress of each in the past.

**Now it is the time to recognize the two
as separate (but complementary) aspects.**



MAJOR PROBLEMS IN CONVENTIONAL MODEL PHYSICS

Major Conceptual Problems

(a) Artificial scale separation by discretization

Discretization introduced for computational purpose separates scales for explicit processes, which can be highly transient, and implicit (parameterized) processes, which can only be near statistical equilibrium. This is especially serious for processes involving clouds.

(b) Incomplete interactions between different physical processes

Due to the modular structure of models, different physical processes (cloud microphysics, radiation, turbulence, etc.) interact mainly through resolved scales, missing most of their small-scale transient interactions.

(c) Non-convergence to "real physics"

Conventional model physics does not converge to local and instantaneous "real physics" as the grid size and time interval approach zero.

The Most Fundamental Technical Problem

Lack of an appropriate cloud-system model (as opposed to a model for individual clouds) to be used as a basic framework

"REAL SOURCES" vs. "REQUIRED SOURCES"

(For prognostic variable ϕ)

Local and instantaneous governing equation:

$$\frac{\partial \phi}{\partial t} = \mathbf{v} \cdot \nabla \phi + \text{"real sources"} \quad (1)$$

Model's governing equation:

$$\frac{\partial \overline{\phi}}{\partial t} = \overline{\mathbf{v} \cdot \nabla \phi} + \text{"required sources"} \quad (2)$$

— : space/time averaging

For (2) to be consistent with $\overline{(1)}$,

"required sources"

$$= \overline{\text{"real sources"}} + \overline{\mathbf{v} \cdot \nabla \phi} - \overline{\mathbf{v} \cdot \nabla \phi}$$

implicit component of advection

(diffusive, penetrative, or . . .)

**Thus, "required physics" is
more than the time/space average of "real physics".**

VS.

"REQUIRED CLOUD-MICROPHYSICAL SOURCES

Similarly to the total physics case,

"required cloud-microphysical sources"


= "real cloud-microphysical sources"

implicit components of

+ advection, turbulence & radiation effects

induced by cloud microphysics


Thus, "required cloud microphysics" is more than the time/space average of "real cloud microphysics".



THE CONVERGENCE PROBLEM

Justification for use of a discrete model relies on the hope that **its solution converges to the true solution as the resolution is refined.**

- "Required sources" for a discrete model to be correct are more than the time/space means of local instantaneous sources ("real sources").
- Naturally, "required sources" converge to the "real sources" as the resolution is refined.
- Conventional model physics is designed and tuned to provide "required sources" for a particular resolution, not those as functions of resolution.



An Example of Non-Convergence


A simple cumulus parameterization

Cumulus warming, $Q_{1c} \sim M_c \partial s / \partial z$

Cumulus drying, $Q_{2c} \sim -M_c L \partial q / \partial z$

Source of moist static energy,

$$Q_{1c} - Q_{2c} \sim M_c \partial h / \partial z$$



Real (i.e., local & instantaneous) cloud-microphysical source of moist static energy

Zero except when ice-phase is involved.

Resolution Dependency of Model Physics

Inferred from Nonhydrostatic Model Experiments

(Jung and Arakawa 2002, submitted to JAS)

- APPLICATIONS OF UCLA/U-UTAH/CSU 2-D CRM to a variety of idealized tropical conditions
- CONTROL RUNS with 2-km horizontal resolution and full physics
- LOW-RESOLUTION RUNS WITH NO OR PARTIAL PHYSICS over a short time interval from selected realizations in CONTROL.
- DIAGNOSIS OF "REQUIRED SOURCES" for the low-resolution model to be correct as far as resolved scales are concerned


Time/Domain Average Profiles of "Real Sources"

Moist Static Energy


No LS Forcing

Strong LS Forcing

Over Land




Over Ocean



Time/Domain Average Profiles of "Real Sources"

Total Water Mixing Ratio




Domain/Ensemble Average Profiles of "Required Cloud-Microphysical Sources"

(From Jung and Arakawa 2002)


Strong LS forcing over land, late afternoon hours

Moist Static Energy

Physics time interval dependence
with 2-km horizontal resolution



Horizontal resolution dependence
with 60 min physics time interval




Domain/Ensemble Average Profiles of "Required Cloud-Microphysical Sources"

(From Jung and Arakawa 2002)

Strong LS forcing over land, late afternoon hours

Total water mixing ratio

Physics time interval dependence
with 2-km horizontal resolution



Horizontal resolution dependence
with 60 min physics time interval

